

Homework 8, due Nov. 30, 2006

1. Problem 13.-3 in book.
2. Consider a system of N **distinguishable** particles, at temperature T , with available energy levels ϵ_1 and $\epsilon_2 > \epsilon_1$ (only two available energy levels).
 - (a) Determine the equilibrium values of the occupation numbers N_1 and N_2 , and the energy U of the system, as a function of temperature.
 - (b) Show that the specific heat is given by

$$C_V = Nk \left(\frac{\Delta}{kT} \right)^2 \frac{e^{-\Delta/kT}}{(1 + e^{-\Delta/kT})^2},$$

where $\Delta = \epsilon_2 - \epsilon_1$. Examine the low temperature and high temperature behavior of C_V/Nk , and sketch it as a function of kT/Δ .

3. Problem 13-8 in book.
 4. Problem 13-9 in book.
 5. Problem 14-6 in book.
 6. Problem 15-2 in book. Note that the result of part (a) can be written as $S = -Nk \sum_j P_j \ln P_j$, where $P_j = N_j/N$ is the probability of occupying level j . In part (c), examine both the $T \rightarrow 0$ and $T \rightarrow \infty$ behavior.
- * . As preparation for the final, also do problems 16-1 and 16-3 in the book. These two problems do not need to be turned in.