

★ **Reading: Luke chapter 9. Tong chapter 4**

• On to fermions! Consider more generally Lorentz transformations. Under Lorentz transformations $x^\mu \rightarrow x^{\mu'} = \Lambda^\mu_{\nu'} x^\nu$, scalar fields transform as $\phi(x) \rightarrow \phi'(x) = \phi(\Lambda^{-1}x)$. Vector fields transform as $A^\mu \rightarrow \Lambda^\mu_{\nu'} A^{\nu'}(\Lambda^{-1}x)$. Generally, $\phi^a \rightarrow D[\Lambda]_b^a \phi^b(\Lambda^{-1}x)$, where $D[\Lambda]$ is a rep of the Lorentz group, $D[\Lambda_1]D[\Lambda_2] = D[\Lambda_1\Lambda_2]$.

Write $D[\Lambda] = \exp(i\frac{1}{2}\Omega_{\mu\nu}\mathcal{M}^{\mu\nu})$, which is a rep if $\mathcal{M}^{\mu\nu}$ satisfies the Lie algebra commutation relation $[\mathcal{M}^{\rho\sigma}, \mathcal{M}^{\mu\nu}] = i\eta^{\sigma\mu}\mathcal{M}^{\rho\nu} \pm 3perms$, where the perms account for $\mathcal{M}^{\mu\nu} = -\mathcal{M}^{\nu\mu}$. E.g. the fundamental rep has $i(\mathcal{M}^{\mu\nu})^{\rho\sigma} = \eta^{\mu\rho}\eta^{\nu\sigma} - \eta^{\mu\sigma}\eta^{\nu\rho}$.

Write the Lorentz transformation generators in terms of rotation, whose generators are the angular momentum \vec{J} , where $J_i = \frac{1}{2}\epsilon_{ijk}M^{jk}$, and boosts, with \vec{K} and $K_i = M^{i,0}$. They are similar, e.g. boosting along the x axis vs rotation around the x axis:

$$\Lambda_{boost} = \begin{pmatrix} \cosh \phi & \sinh \phi & & \\ \sinh \phi & \cosh \phi & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \quad \Lambda_{rotate} = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \cos \theta & -\sin \theta \\ & & \sin \theta & \cos \theta \end{pmatrix}.$$

So define $\vec{N}^\pm \equiv \frac{1}{2}(\vec{J} \pm i\vec{K})$. Then the Lorentz algebra becomes simply $[N_i^\pm, N_k^\pm] = i\epsilon_{ijk}N_k^\pm$, and $[N^\pm, N_j^\mp] = 0$, i.e. two copies of the familiar rotation commutation relations. The reps are then labeled by (n_L, n_R) , where n_L and n_R are non-negative half-integers, like the angular momentum j . Note that parity exchanges $\vec{N} \leftrightarrow \vec{N}^\dagger$, so it exchanges the above left and right, hence their names. The angular momentum $\vec{J} = \vec{N} + \vec{N}^\dagger$, so j runs from $|n_L - n_R|$ to $n_L + n_R$. The scalar rep is $(0, 0)$, the vector rep is $(1/2, 1/2)$.¹ The basic spinor reps are $(1/2, 0)$ and $(0, 1/2)$, denoted u_\pm ; these are called left and right handed Weyl spinors. They both have $D = e^{-i\vec{\sigma} \cdot \hat{e}\theta/2}$ for a rotation by θ around the \hat{e} axis, but they have $D_\pm = e^{\pm\vec{\sigma} \cdot \hat{e}\phi/2}$ for a boost along the \hat{e} axis, where $v = \tanh \phi$. These 2-component Weyl spinor representations individually play an important role in non-parity invariant theories, like the weak interactions. Parity $((t, \vec{x}) \rightarrow (t, -\vec{x}))$ exchanges them. So, in

¹ Consider $\sigma^\mu = (1, \sigma^i)$, where each entry is a 2×2 matrix. Now form $X = x^\mu \sigma^\mu u$. Lorentz transformations act as $X \rightarrow X' = DXD^\dagger$, where $D \in SL(2, C)$. Here $D = e^{-i\vec{\sigma} \cdot \hat{e}\theta/2}$ for a rotation by θ around the \hat{e} axis, and $D_\pm = e^{\pm\vec{\sigma} \cdot \hat{e}\phi/2}$ for a boost along the \hat{e} axis, where $v = \tanh \phi$. This illustrates the statement that the vector representation of the Lorentz group is $D^{(1/2, 1/2)}$.

parity invariant theories, like QED, they are combined into a 4-component Dirac spinor, $(1/2, 0) + (0, 1/2)$:

$$\psi = \begin{pmatrix} u_+ \\ u_- \end{pmatrix}.$$

The 4-component spinor rep starts with the clifford algebra $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}\mathbf{1}$, e.g.

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}.$$

There are other choices of reps of the clifford algebra.

$S^{\mu\nu} = \frac{1}{4}[\gamma^\mu, \gamma^\nu] = \frac{1}{2}\gamma^\mu\gamma^\nu - \frac{1}{2}\eta^{\mu\nu}$, satisfies the Lorentz Lie algebra relation. Under a rotation, $S^{ij} = -\frac{i}{2}\epsilon_{ijk} \begin{pmatrix} \sigma^k & 0 \\ 0 & \sigma^k \end{pmatrix}$, so taking $\Omega_{ij} = -\epsilon_{ijk}\varphi^k$ get under rotations

$$S[\vec{\varphi}] = \begin{pmatrix} e^{i\vec{\varphi}\cdot\vec{\sigma}/2} & 0 \\ 0 & e^{i\vec{\varphi}\cdot\vec{\sigma}/2} \end{pmatrix}.$$

Under boosts, $\Omega_{i,0} = \phi_i$,

$$S[\Lambda] = \begin{pmatrix} e^{\vec{\phi}\cdot\vec{\sigma}/2} & 0 \\ 0 & e^{-\vec{\phi}\cdot\vec{\sigma}/2} \end{pmatrix}.$$

This exhibits the 2-component reps that we described above.

Under Lorentz transformations, spinors transform as $\psi(x) \rightarrow S[\Lambda]\psi(\Lambda^{-1}x)$, and $\psi^\dagger(x) \rightarrow \psi^\dagger(\Lambda^{-1}x)S[\Lambda]^\dagger$. Note that $S[\Lambda]^\dagger S[\Lambda] \neq 1$, but $S[\Lambda]^\dagger = \gamma^0 S[\Lambda]^{-1} \gamma^0$. So define $\bar{\psi}(x) \equiv \psi^\dagger \gamma^0$ and note that $\bar{\psi}\psi$ transforms as a scalar, and $\bar{\psi}\gamma^\mu\psi$ transforms as a Lorentz 4-vector.

For 2-component spinors, $u_-^\dagger \sigma^\mu u_-$ and $u_+^\dagger \bar{\sigma}^\mu u_+$ transform like vectors, where $\sigma^\mu = (1, \sigma^i)$ and $\bar{\sigma}^\mu = (1, -\sigma^i)$. Here are two Lorentz scalars (exchanged under parity): $u_\pm^\dagger u_\mp$.

$\gamma^5 \equiv -i\gamma^0\gamma^1\gamma^2\gamma^3$, anticommutes with all other γ^μ and $(\gamma^5)^2 = 1$. In our above representation of the gamma matrices, $\gamma_5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, so $P_\pm = \frac{1}{2}(1 \pm \gamma^5)$ are projection operators, projecting on to u_\pm .

- The Dirac action:

$$\begin{aligned} S &= \int d^4x \bar{\psi}(x) (i\gamma^\mu \partial_\mu - m) \psi(x) \\ &= \int d^4x (u_+^\dagger i\sigma^\mu \partial_\mu u_+ + u_-^\dagger i\bar{\sigma}^\mu \partial_\mu u_- - m(u_+^\dagger u_- + u_-^\dagger u_+)). \end{aligned}$$