

5/6/13 Lecture outline

★ Reading: Zwiebach chapter 7.

● Recall from last time:

$$\mathcal{L}_{NG} = -\frac{T_0}{c} \sqrt{(\dot{X} \cdot X')^2 - (\dot{X})^2 (X')^2},$$

and we have

$$\mathcal{P}_\mu^\tau = \frac{\partial \mathcal{L}}{\partial \dot{X}^\mu} = -\frac{T_0}{c} \frac{(\dot{X} \cdot X') X'_\mu - (X')^2 \dot{X}_\mu}{\sqrt{(\dot{X} \cdot X')^2 - (\dot{X})^2 (X')^2}},$$

and

$$\mathcal{P}_\mu^\sigma = \frac{\partial \mathcal{L}}{\partial X^{\mu'}} = -\frac{T_0}{c} \frac{(\dot{X} \cdot X') \dot{X}_\mu - (\dot{X})^2 X'_\mu}{\sqrt{(\dot{X} \cdot X')^2 - (\dot{X})^2 (X')^2}}.$$

The condition $\delta S = 0$ gives the Euler-Lagrange equations

$$\frac{\partial \mathcal{P}_\mu^\tau}{\partial \tau} + \frac{\partial \mathcal{P}_\mu^\sigma}{\partial \sigma} = 0.$$

Exploit $(\tau, \sigma) \rightarrow (\tau', \sigma')$ reparameterization invariance to pick useful “gauges”, to simplify the above equations. We will discuss choices such that we can impose constraints

$$\dot{X} \cdot X' = 0 \quad \dot{X}^2 + X'^2 = 0. \quad (1)$$

In this case, we have

$$\mathcal{P}^{\tau\mu} = \frac{1}{2\pi\alpha'} \dot{X}^\mu \quad \mathcal{P}^{\sigma\mu} = -\frac{1}{2\pi\alpha'} X^{\mu'}, \quad (2)$$

and then the EOM is simply a wave equation:

$$(\partial_\tau^2 - \partial_\sigma^2) X^\mu = 0. \quad (3)$$

Step 1 (last time): Static gauge: pick $\tau = t$. Verify sign inside $\sqrt{\cdot}$ in this case: $X^{\mu'} = (0, \vec{X}')$, $\dot{X}^\mu = (c, \dot{\vec{X}})$, take e.g. $\dot{\vec{X}} = 0$ to get $\sqrt{\cdot} = c|\vec{X}'|$. Express S in terms of $\vec{v}_\perp = \partial_t \vec{X} - (\partial_t \vec{X} \cdot \partial_s \vec{X}) \partial_s \vec{X}$ (with $ds \equiv |d\vec{X}|_{t=const} = |\partial_\sigma \vec{X}| |d\sigma|$), show $(\dot{X} \cdot X')^2 - \dot{X}^2 (X')^2 = (\frac{ds}{d\sigma})^2 (c^2 - v_\perp^2)$, to get $L = -T_0 \int ds \sqrt{1 - v_\perp^2/c^2}$. Also get

$$\mathcal{P}^{\sigma\mu} = -\frac{T_0}{c^2} \frac{(\partial_s \vec{X} \cdot \partial_t \vec{X}) \dot{X}^\mu + (c^2 - (\partial_t \vec{X})^2) \partial_s X^\mu}{\sqrt{1 - v_\perp^2/c^2}},$$

$$\mathcal{P}^{\tau\mu} = \frac{T_0}{c^2} \frac{ds}{d\sigma} \frac{\dot{X}^\mu - (\partial_s \vec{X} \cdot \partial_t \vec{X}) \partial_s X^\mu}{\sqrt{1 - v_\perp^2/c^2}}.$$

- Free, Neuman BCs, P_μ^σ for the $\mu = 0$ component implies that endpoints move transversely, $\partial_s \vec{X} \cdot \partial_t \vec{X} = 0$, so $\vec{v}_\perp = \vec{v}$. The condition $\vec{P}^\sigma = 0$ at the endpoints implies that the speed of light, $v = c$, for the free (Neuman) BCs.

- Step 2: can choose σ such that $\partial_\sigma \vec{X} \cdot \partial_t \vec{X} = 0$ along entire string (we saw it above for the endpoints). This gives $\vec{v}_\perp = \vec{v} \equiv \dot{\vec{X}}$ along the entire string. Then $\mathcal{P}^{\tau\mu} = \frac{T_0}{c^2} \frac{ds}{d\sigma} \gamma \partial_t X^\mu$ and $\mathcal{P}^{\sigma\mu} = -T_0 \gamma^{-1} \partial_s X^\mu$, with $\gamma \equiv 1/\sqrt{1 - v_\perp^2/c^2}$.

Now consider the $\mu = 0$ component of the EOM: $\partial_t \mathcal{P}^{\tau\mu} = -\partial_\sigma \mathcal{P}^{\sigma\mu}$, which for $\mu = 0$ gives that $(T_0/c) \frac{ds}{d\sigma} \gamma$ is a constant of the motion. Indeed this is proportional to the energy of an element of string. Now the space components of the EOM can be written as $\mu_{eff} \partial_t \vec{v}_\perp = \partial_s (T_{eff} \partial_s \vec{X})$, with $T_{eff} = T_0/\gamma$ and $\mu_{eff} = T_0 \gamma/c^2$.

- Now note that since $\frac{ds}{d\sigma} \gamma$ is a constant, we can set it equal to 1. This can be written as the constraint: $(\partial_\sigma \vec{X})^2 + (\partial_{X_0} \vec{X})^2 = 1$.

- Summary: choose σ parameterization such that

$$\partial_\sigma \vec{X} \cdot \partial_\tau \vec{X} = 0 \quad \text{and} \quad d\sigma = \frac{ds}{\sqrt{1 - v_\perp^2/c^2}} = \frac{dE}{T_0}.$$

(Using $H = \int T_0 ds / \sqrt{1 - v_\perp^2/c^2}$ and $\partial_t (ds / \sqrt{1 - v_\perp^2/c^2}) = 0$.) The last equation above is equivalent to $(\partial_\sigma \vec{X})^2 + c^{-2} (\partial_t \vec{X})^2 = 1$. With this worldsheet gauge choice,

$$\mathcal{P}^{\tau\mu} = \frac{T_0}{c^2} \partial_t X^\mu = \frac{T_0}{c^2} (c, \vec{v}_\perp), \quad \mathcal{P}^{\sigma,\mu} = -T_0 \partial_\sigma X^\mu = (0, -T_0 \partial_\sigma \vec{X}).$$

We can write this as

$$\mathcal{P}^{\tau\mu} = \frac{1}{2\pi\alpha'} \dot{X}^\mu \quad \mathcal{P}^{\sigma\mu} = -\frac{c^2}{2\pi\alpha'} X^{\mu'}, \quad (4)$$

and then the EOM is simply a linear wave equation, and we also need to impose the constraints:

$$(\partial_\tau^2 - c^2 \partial_\sigma^2) X^\mu = 0, \quad (\dot{X} \pm X')^2 = 0. \quad (5)$$

- Solution of the EOM for open string with free BCs at each end: imposing first at $\sigma = 0$ gives $\vec{X}(t, \sigma) = \frac{1}{2} (\vec{F}(ct + \sigma) + \vec{F}(ct - \sigma))$ where the open string has $\sigma \in [0, \sigma_1]$ and (1) implies that $|\frac{d\vec{F}(u)}{du}|^2 = 1$, and $\vec{X}'|_{ends} = 0$ implies $\vec{F}(u + 2\sigma_1) = \vec{F}(u) + 2\sigma_1 \vec{v}_0/c$. Note $\vec{F}(u)$ is the position of the $\sigma = 0$ end at time u/c . Then show that \vec{v}_0 is the average velocity of any point σ on the string over time interval $2\sigma_1/c$. Observing motion of $\sigma = 0$ end over that Δt , together with E , gives motion of string for all t . Example from book: $\vec{X}(t, \sigma = 0) = \frac{\ell}{2} (\cos \omega t, \sin \omega t)$. Find $\vec{F}(u) = \frac{\sigma_1}{\pi} (\cos \pi u / \sigma_1, \sin \pi u / \sigma_1)$, with $\vec{v}_0 = 0$. $|\frac{d\vec{F}}{du}|^2 = 1$ gives $\ell = 2c/\omega = 2E/\pi T_0$. Finally, $\vec{X}(t, \sigma) = \frac{\sigma_1}{\pi} \cos(\pi \sigma / \sigma_1) (\cos(\pi ct / \sigma_1), \sin(\pi ct / \sigma_1))$.