

2/12/10 Lecture 12 outline

- Last time, mentioned renormalized and bare Greens functions. Recall that  $\Phi_B \equiv Z_\phi^{1/2} \phi$ , and the definition of the 1PI Green's functions  $\tilde{\Gamma}^{(n)}$ , and in particular that they have all  $n$  external propagators amputated. It then follows that

$$\tilde{\Gamma}_B^{(n)}(p_1, \dots, p_n; \lambda_B, m_B, \epsilon) = Z_\phi^{-n/2} \tilde{\Gamma}_R^{(n)}(p_1, \dots, p_n; \lambda_R, m_R, \mu, \epsilon).$$

For fixed physics, the LHS is some fixed quantity. The RHS depends on the renormalization point  $\mu$  and the scheme. The LHS does not! This leads to what is known as the renormalization group equations, which state how the renormalized quantities must vary with  $\mu$ . Rewrite above as

$$Z_\phi^{n/2} \tilde{\Gamma}_B^{(n)}(p_1, \dots, p_n; \lambda_B, m_B, \epsilon) = \tilde{\Gamma}_R^{(n)}(p_1, \dots, p_n; \lambda_R, m_R, \mu, \epsilon).$$

Now the RHS is finite, so the LHS must be too. So we can take  $\epsilon \rightarrow 0$  without a problem.

- Before getting into the renormalization group, let's take a little detour. Recall that

$$\int d^4x e^{ipx} \langle \Omega | T \phi(x) \phi(0) | \Omega \rangle = \frac{i}{p^2 - m^2 - \Pi'(p^2) + i\epsilon}.$$

Here  $|\Omega\rangle$  is the full, interacting vacuum and  $\phi$  are the full (Heisenberg picture) operators. Now insert a complete set of states,

$$\mathbf{1} = |\Omega\rangle\langle\Omega| + \sum_\lambda \int \frac{d^3p}{(2\pi)^2} \frac{1}{2E_p(\lambda)} |\lambda_p\rangle\langle\lambda_p|$$

where  $\lambda$  are all eigenstates of the full  $H$ , and  $\lambda_p$  is a boosted version, to give an eigenstate of  $\vec{P}$ , with spatial momentum  $\vec{p}$ . Now use  $\langle \Omega | \phi(x) | \lambda_p \rangle = \langle \Omega | \phi(x) | \lambda_0 \rangle e^{-ipx}$  (where  $p^0 = E_p \equiv \sqrt{|\vec{p}|^2 + m_\lambda^2}$ ). Use now  $\int \frac{d^3\vec{p}}{(2\pi)^3 2E_p} \rightarrow \int \frac{d^4p}{(2\pi)^4} \frac{i}{p^2 - m_\lambda^2 + i\epsilon}$  to show that the LHS can be written as

$$\int d^4x e^{ipx} \langle \Omega | T \phi(x) \phi(0) | \Omega \rangle = \int_0^\infty \frac{dM^2}{2\pi} \rho(M^2) \frac{i}{p^2 - M^2 + i\epsilon},$$

where

$$\rho(M^2) = \sum_\lambda 2\pi \delta(M^2 - m_\lambda^2) |\langle \Omega | \phi(0) | \lambda \rangle|^2 > 0$$

is the Kallen-Lehmann spectral density. Find  $\rho(M^2) = 2\pi \delta(M^2 - m^2) Z$  for  $M^2 \ll 4m^2$ . For  $M^2$  slightly below  $4m^2$  there are new delta functions, at the bound states. Starting at  $4m^2$ ,  $\rho(M^2)$  is some positive function. This implies that

$$\frac{i}{p^2 - m^2 - \Pi'(p^2) + i\epsilon} = \frac{iZ}{p^2 - m^2 + i\epsilon} + \int_{\sim 4m^2}^\infty \frac{dM^2}{2\pi} \rho(M^2) \frac{i}{p^2 - M^2 + i\epsilon}.$$

The LHS has a simple pole, with residue  $iZ$ , at  $p^2 - m^2$ . Here  $Z = |\langle \lambda_0 | \phi(0) | \Omega \rangle|^2$  is the probability for  $\phi(0)$  to create the lowest energy 1-particle state from the vacuum. Then there can be a few more simple poles, for  $p^2$  slightly below  $4m^2$ .

Starting at  $p^2 = 4m^2$ , there is a branch cut, corresponding to producing two more free particles. Note  $\mathcal{M}(s) = \mathcal{M}(s^*)^*$  implies that the real part of  $\mathcal{M}$  is continuous across the cut, but the imaginary part can be discontinuous:  $Im\mathcal{M}(s + i\epsilon) = -Im\mathcal{M}(s - i\epsilon)$ . We'll return to this shortly.

The above equality, back in position space and taking  $\partial/\partial t$ , leads to the equal time commutators,  $[\phi(\vec{x}, t), \dot{\phi}(\vec{y}, t)] = i\delta^{(3)}(\vec{x} - \vec{y})$ , matching the coefficient of the delta function on the two sides of the resulting equation gives

$$1 = Z + \int_{\sim 4m^2}^{\infty} \frac{dM^2}{2\pi} \rho(M^2) \geq Z.$$

Implies that  $0 \leq Z \leq 1$ , with  $Z = 1$  iff the theory is a free field theory. Fits with what we found before,

$$\delta_Z^{(2)} = -\frac{\lambda^2}{12(16\pi^2)^2} \frac{1}{\epsilon},$$

so negative (for  $\epsilon > 0$ ).

- Recall LSZ (Lehmann, Symanzik, Zimmermann '55) from last quarter, now noting that there are  $Z$  factors. Let's just state the result: the S-matrix element for  $m$  incoming particles and  $n$  outgoing ones is given by

$$\langle \mathbf{p}_1 \dots \mathbf{p}_n | S | \mathbf{k}_1 \dots \mathbf{k}_m \rangle = \lim_{o.s.} \prod_{i=1}^n (p_i^2 - m_i^2) Z_i^{-1/2} \prod_{j=1}^m (k_j^2 - m_j^2) Z_j^{-1/2} \tilde{G}^{n+m}(-p_i, k_i).$$

Here  $\tilde{G}^{n+m}$  is the full  $n + m$  point Green's function, including disconnected diagrams etc. The limit is where we take the external particles on shell. In this limit, the  $p_i^2 - m_i^2$  and  $k_j^2 - m_j^2$  prefactors all go to zero. These zeros kill everything on the RHS except for the connected contributions to  $\tilde{G}$ . Accounting for the fact that we amputate the external propagators, which go like  $iZ_i(p_i^2 - m_i^2)^{-1}$ , the above becomes

$$\langle \mathbf{p}_1 \dots \mathbf{p}_n | S | \mathbf{k}_1 \dots \mathbf{k}_m \rangle = Z^{(n+m)/2} \tilde{G}_{conn,B}^{n+m}(-p_i, k_i) = \tilde{G}_{conn,R}^{n+m}(-p_i, k_j)$$